

Visibility and the Birch and Swinnerton-Dyer conjecture for analytic rank one

Amod Agashe *

October 14, 2008

Abstract

Let E be an optimal elliptic curve over $\mathbf{Q}$ of conductor N having analytic rank one, i.e., such that the L -function $L_E(s)$ of E vanishes to order one at $s = 1$. Let K be a quadratic imaginary field in which all the primes dividing N split and such that the L -function of E over K vanishes to order one at $s = 1$. Suppose there is another optimal elliptic curve over $\mathbf{Q}$ of the same conductor N whose Mordell-Weil rank is greater than one and whose associated newform is congruent to the newform associated to E modulo an integer r . The theory of visibility then shows that under certain additional hypotheses, r divides the order of the Shafarevich-Tate group of E over K . We show that under somewhat similar hypotheses, r also divides the Birch and Swinnerton-Dyer *conjectural* order of the Shafarevich-Tate group of E over K , which provides new theoretical evidence for the second part of the Birch and Swinnerton-Dyer conjecture in the analytic rank one case.

1 Introduction and results

Mazur introduced the notion of visibility in order to better understand geometrically the elements of the Shafarevich-Tate group of an abelian variety. The corresponding theory, which we call the theory of visibility, can often be used to show the existence of non-trivial elements of the Shafarevich-Tate group of abelian varieties and motives (e.g., see [CM00], [AS02b], [DSW03]). The second part of the Birch and Swinnerton-Dyer conjecture gives a formula for the order of the Shafarevich-Tate group, and one might wonder how

*This material is based upon work supported by the National Science Foundation under Grant No. 0603668.

much of this conjectural order, when non-trivial, can be explained by the theory of visibility. This issue has been investigated computationally (e.g., see [CM00], [AS05]) and theoretically (e.g., [Aga07]) when the concerned abelian variety has analytic rank zero, but not for any higher analytic rank (where, by the *analytic rank* of an abelian variety, we mean the order of vanishing of the L -function of the abelian variety at $s = 1$). In this article, we take the first step in investigating the issue for higher analytic rank, by showing that for elliptic curves of analytic rank one, in certain situations where the theory of visibility implies that the actual Shafarevich-Tate group is non-trivial, the Birch and Swinnerton-Dyer conjectural order of the Shafarevich-Tate group is non-trivial as well. This provides new theoretical evidence for the second part of the Birch and Swinnerton-Dyer conjecture in the analytic rank one case, and links the theory of visibility to the Birch and Swinnerton-Dyer conjecture for the first time for any elliptic curve of analytic rank greater than zero.

We now state our results more precisely. Let N be a positive integer. Let $X_0(N)$ be the modular curve over $\mathbf{Q}$ associated to $\Gamma_0(N)$, and let $J = J_0(N)$ denote the Jacobian of $X_0(N)$, which is an abelian variety over $\mathbf{Q}$. Let $\mathbf{T}$ denote the Hecke algebra, which is the subring of endomorphisms of $J_0(N)$ generated by the Hecke operators (usually denoted T_ℓ for $\ell \nmid N$ and U_p for $p \mid N$). If f is a newform of weight 2 on $\Gamma_0(N)$, then let $I_f = \text{Ann}_{\mathbf{T}} f$ and let A_f denote the associated *newform quotient* $J/I_f J$, which is an abelian variety over $\mathbf{Q}$. Let π denote the quotient map $J \rightarrow J/I_f J = A_f$. By the *analytic rank* of f , we mean the order of vanishing at $s = 1$ of $L(f, s)$. The analytic rank of A_f is then the analytic rank of f times the dimension of A_f . Now suppose that the newform f has integer Fourier coefficients. Then A_f is an elliptic curve, and we denote it by E instead. Since E has dimension one, it has analytic rank one.

Let K be a quadratic imaginary field of discriminant not equal to -3 or -4 , and such that all primes dividing N split in K . Choose an ideal $\mathcal{N}$ of the ring of integers $\mathcal{O}_K$ of K such that $\mathcal{O}_K/\mathcal{N} \cong \mathbf{Z}/N\mathbf{Z}$. Then the complex tori $\mathbf{C}/\mathcal{O}_K$ and $\mathbf{C}/\mathcal{N}^{-1}$ define elliptic curves related by a cyclic N -isogeny, and thus give a complex valued point x of $X_0(N)$. This point, called a Heegner point, is defined over the Hilbert class field H of K . Let $P \in J(K)$ be the class of the divisor $\sum_{\sigma \in \text{Gal}(H/K)} ((x) - (\infty))^\sigma$, where H is the Hilbert class field of K .

By [Wal85], we may choose K so that $L(E/K, s)$ vanishes to order one at $s = 1$. Hence, by [GZ86, §V.2:(2.1)], $\pi(P)$ has infinite order, and by work of Kolyvagin, $E(K)$ has rank one and the order of the Shafarevich-Tate group $\text{III}(E/K)$ of E over K is finite (e.g., see [Kol90, Thm. A] or [Gro91,

Thm. 1.3]). In particular, the index $[E(K) : \mathbf{Z}\pi(P)]$ is finite. By [GZ86, §V.2:(2.2)] (or see [Gro91, Conj. 1.2]), the second part of the Birch and Swinnerton-Dyer (BSD) conjecture becomes:

Conjecture 1.1 (Birch and Swinnerton-Dyer, Gross-Zagier).

$$|E(K)/\mathbf{Z}\pi(P)| \stackrel{?}{=} c_E \cdot \prod_{p|N} c_p(E) \cdot |\text{III}(E/K)|^{1/2}, \quad (1)$$

where c_E is the Manin constant of E , $c_p(E)$ denotes the arithmetic component group of E at the prime p , and the question mark above the equality sign emphasizes that this equality is conjectural.

Note that the Manin constant c_E is conjectured to be one, and one knows that if p is a prime such that $p^2 \nmid 4N$, then p does not divide c_E (by [Maz78, Cor. 4.1] and [AU96, Thm. A]).

Now suppose that f is congruent to another newform g with integer Fourier coefficients, whose associated elliptic curve F has Mordell-Weil rank over $\mathbf{Q}$ bigger than one. Then the theory of visibility (e.g., as in [CM00]) often shows that such congruences divide the order of $\text{III}(E/K)$, as we now indicate.

We say that a maximal ideal $\mathfrak{m}$ of $\mathbf{T}$ satisfies *multiplicity one* if $J_0(N)[\mathfrak{m}]$ is two dimensional over $\mathbf{T}/\mathfrak{m}$. This is known to hold in several situations, in particular when the following conditions hold simultaneously (e.g., see [Wil95, Thm. 2.1(ii)] along with [ARS07]): $p \neq 2$, $p^2 \nmid N$, the canonical semi-simple representation $\rho_{\mathfrak{m}}$ associated to $\mathfrak{m}$ (see, e.g., [Rib90, Prop. 5.1] for the definition of $\rho_{\mathfrak{m}}$) is irreducible, and $\mathfrak{m}$ arises as a pullback from $\mathbf{T}/I_h$ for some newform h .

Lemma 1.2. *Let r be an integer such that:*

- (a) *f and g are congruent modulo r and for every prime p that divides r , f and g are not congruent modulo a power of p greater than $p^{\text{ord}_p r}$.*
- (b) *if $\mathfrak{m}$ is a maximal ideal of $\mathbf{T}$ such that the residue characteristic of $\mathfrak{m}$ divides r and $\mathfrak{m}$ is in the support of $J_0(N)[I_f]$ or $J_0(N)[I_g]$, then $\mathfrak{m}$ satisfies multiplicity one.*

Then $E^\vee[r] = F^\vee[r]$, and both are direct summands of $E^\vee \cap F^\vee$ as $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ -modules.

Proof. By [Eme03, Cor. 2.5], if $\mathfrak{m}$ satisfies multiplicity one and I is any saturated ideal of $\mathbf{T}$, then the $\mathfrak{m}$ -adic completion of the group of connected components of $J_0(N)[I]$ is trivial. If $L \rightarrow M$ is a homomorphism of two $\mathbf{T}$ -modules, then we say that $L = M$ *away from* a given set of maximal ideals

if the induced map on the $\mathfrak{m}$ -adic completions is an isomorphism for all maximal ideals $\mathfrak{m}$ that are not in the prescribed set. Thus, the inclusions $E^\vee \subseteq J_0(N)[I_f]$ and $F^\vee \subseteq J_0(N)[I_g]$ are equalities away from maximal ideals that do not satisfy multiplicity one. Hence $E^\vee \cap F^\vee \subseteq J_0(N)[I_f + I_g]$ is an equality away from maximal ideals that do not satisfy multiplicity one. But $E^\vee \cap F^\vee \subseteq E^\vee[I_f + I_g] \subseteq J_0(N)[I_f + I_g]$. Hence if m denotes the largest integer such that f and g are congruent modulo m , then $E^\vee \cap F^\vee \subseteq E^\vee[I_f + I_g] = E^\vee[m]$ is an equality away from maximal ideals that do not satisfy multiplicity one. Similarly $E^\vee \cap F^\vee \subseteq F^\vee[I_f + I_g] = F^\vee[m]$ is an equality away from maximal ideals that do not satisfy multiplicity one. From conditions (a) and (b) on r , it follows that for every prime p that divides r , $(E \cap F)[p^\infty] = E^\vee[p^\infty] = F^\vee[p^\infty]$. Thus $E^\vee[r]$ and $F^\vee[r]$ are identical and are direct summands of $E^\vee \cap F^\vee$ as $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ -modules. $\square$

Proposition 1.3. (i) *Let r be the largest integer such that f and g are congruent modulo r and all maximal ideals $\mathfrak{m}$ of $\mathbf{T}$ such that the residue characteristic of $\mathfrak{m}$ divides r and $\mathfrak{m}$ is in the support of $J_0(N)[I_f]$ or $J_0(N)[I_g]$ satisfy multiplicity one. Suppose that r is coprime to*

$$N \cdot |(J_0(N)/F^\vee)(K)_{\text{tor}}| \cdot |F(K)_{\text{tor}}| \cdot \prod_{p|N} c_p(F).$$

Then r divides $\prod_{p|N} c_p(E) \cdot |\text{III}(E/K)|^{1/2}$, which in turn divides the right hand side of the Birch and Swinnerton-Dyer conjectural formula (1).

(ii) *Suppose that f is congruent to g modulo an odd prime q such that $E[q]$ and $F[q]$ are irreducible and q does not divide*

$$N \cdot |(J_0(N)/F^\vee)(K)_{\text{tor}}| \cdot |F(K)_{\text{tor}}|.$$

Also, assume that f is not congruent modulo q to a newform of a level dividing N/p for some prime p that divides N (for Fourier coefficients of index coprime to Nq), and either $q \nmid N$ or for all primes p that divide N , $q \nmid (p-1)$. Then q divides $|\text{III}(E/K)|$.

Proof. Both results follow essentially from Theorem 3.1 of [AS02b]. For the first part, take $A = E^\vee$, $B = F^\vee$, and $n = r$ in [AS02b, Thm. 3.1], and note that $F^\vee[r] \subseteq E^\vee$ by Lemma 1.2 and that the rank of $E^\vee(K)$ is less than the rank of $F^\vee(K)$. For the second part, take $A = E^\vee$, $B = F^\vee$, and $n = q$ in [AS02b, Thm. 3.1], and note that the congruence of f and g modulo q forces $F^\vee[q] = E^\vee[q]$ by [Rib90, Thm. 5.2] (cf. [CM00, p. 20]), and that the hypotheses imply that q does not divide $c_p(E)$ or $c_p(F)$ for any prime p that divides N , as we now indicate. By [Eme03, Prop. 4.2], if

q divides $c_p(E)$ for some prime p that divides N , then for some maximal ideal $\mathfrak{m}$ of $\mathbf{T}$ having characteristic q and containing I_f , either $\rho_{\mathfrak{m}}$ is finite or reducible. Since $E[q]$ is irreducible, this can happen only if $\rho_{\mathfrak{m}}$ is finite. But this is not possible by [Rib90, Thm. 1.1], in view of the hypothesis that f is not congruent modulo q to a newform of a level dividing N/p for some prime p that divides N (for Fourier coefficients of index coprime to Nq), and either $q \nmid N$ or for all primes p that divide N , $q \nmid (p-1)$. Thus q does not divide $c_p(E)$ for any prime p that divides N . Similarly, q does not divide $c_p(F)$ for any prime p that divides N , considering that the hypothesis that f is not congruent modulo q to a newform of a level dividing N/p for some prime p that divides N (for Fourier coefficients of index coprime to Nq) applies to g as well, since g is congruent to f modulo q . This finishes the proof of the proposition. $\square$

One might wonder how often it happens in numerical data that visibility explains the Shafarevich-Tate group of an elliptic curve of analytic rank one. Since it is difficult to compute the actual order of the Shafarevich-Tate group, we looked at the Birch and Swinnerton-Dyer conjectural orders in Cremona's online "Elliptic curve data" [Cre]. For levels up to 30000, we found only one optimal elliptic curve of Mordell-Weil rank one for which the conjectural order of the Shafarevich-Tate group was divisible by an odd prime: the curve with label 28042A, for which the conjectural order of the Shafarevich-Tate group is 9. At the same level, the curve 28042B has Mordell-Weil rank 3 and the newforms corresponding to 28042A and 28042B have Fourier coefficients that are congruent modulo 3 for every prime index up to 100. While this is not enough to conclude that the newforms are congruent modulo 3 for all Fourier coefficients (cf. [AS02a]), it is quite likely that this is true and that this congruence explains the non-trivial Shafarevich-Tate group, although we have not checked the details (in particular whether the hypotheses of Proposition 1.3 are satisfied), since our goal in this paper is to prove *theoretical* results. It would be interesting to do systematic computations to see how much of the Birch and Swinnerton-Dyer conjectural order of the Shafarevich-Tate group is explained by visibility for elliptic curves of analytic rank one (similar to the computations in [AS05] for the analytic rank zero case).

In any case, when the theory of visibility does imply that the Shafarevich-Tate group is non-trivial, this should be reflected in its Birch and Swinnerton-Dyer conjectural order. In our situation, since Proposition 1.3 shows that under certain hypotheses, certain integers of congruence divide the right side of the Birch and Swinnerton-Dyer conjectural formula (1), these inte-

gers should also divide the left side of (1), if the second part of the Birch and Swinnerton-Dyer conjecture is true. We have the following result in this direction.

Theorem 1.4. *Recall that E and F are optimal elliptic curves over $\mathbf{Q}$ associated to newforms f and g respectively, with E of analytic rank one and F having Mordell-Weil rank more than one. Let r be an integer such that:*

- (a) *f and g are congruent modulo r and for every prime p that divides r , f and g are not congruent modulo a power of p greater than $p^{\text{ord}_p r}$.*
- (b) *all maximal ideals $\mathfrak{m}$ of $\mathbf{T}$ whose residue characteristic divides r and that are in the support of $J_0(N)[I_f]$ or of $J_0(N)[I_g]$ satisfy multiplicity one.*
- (c) *r is coprime to the order of the torsion subgroup of the projection of $\mathbf{TP}$ in $J_0(N)/(I_f \cap I_g)J_0(N)$.*

Then r divides $|E(K)/\mathbf{Z}\pi(P)|$, which is the left hand side of the Birch and Swinnerton-Dyer conjectural formula (1).

We will prove this theorem in Section 2.

Corollary 1.5. *Suppose f and g are congruent modulo an odd prime q such that $q^2 \nmid N$, $E[q]$ and $F[q]$ are irreducible, and q does not divide $|J_0(N)(K)_{\text{tor}}|$. Then q divides $|E(K)/\mathbf{Z}\pi(P)|$. If moreover, f is not congruent modulo q to a newform of a level dividing N/p for some prime p that divides N (for Fourier coefficients of index coprime to Nq), and either $q \nmid N$ or for all primes p that divide N , $q \nmid (p-1)$, then q divides the Birch and Swinnerton-Dyer conjectural order of $\text{III}(E/K)$.*

Proof. Let r denote the highest power of q modulo which f and g are congruent. Thus condition (a) on r in Theorem 1.4 is satisfied. By the discussion just before Lemma 1.2, the hypotheses that q is odd, $q^2 \nmid N$, and $E[q]$ and $F[q]$ are irreducible imply that r satisfies condition (b). The hypothesis that q does not divide $|J_0(N)(K)_{\text{tor}}|$ implies that condition (c) is satisfied. Hence by Theorem 1.4, q divides $|E(K)/\mathbf{Z}\pi(P)|$. As explained in the proof of Proposition 1.3, the hypotheses imply that q does not divide $c_p(E)$ for any prime p . Also, by [Maz78, Cor. 4.1], q does not divide the Manin constant c_E . Hence, by (1), q divides the Birch and Swinnerton-Dyer conjectural order of $\text{III}(E/K)$. $\square$

In view of Proposition 1.3, Theorem 1.4 and Corollary 1.5 provide theoretical evidence towards the Birch and Swinnerton-Dyer conjectural formula (1). We remark that Theorem 1.4 is to be compared to part (i) of Proposition 1.3 and Corollary 1.5 to part (ii) of Proposition 1.3. Regarding

the hypothesis in Corollary 1.5 that q does not divide $|J_0(N)(K)_{\text{tor}}|$, we do not know of any results that would give some criteria on q which would imply that this hypothesis holds (unlike the similar situation over $\mathbf{Q}$, where at least for prime N , we know by [Maz77, Thm (1)] that $|J_0(N)(\mathbf{Q})_{\text{tor}}|$ equals the numerator of $\frac{N-1}{12}$). As in Theorem 1.4, we could have replaced this hypothesis by the requirement that p is coprime to the order of the torsion subgroup of the projection of $\mathbf{TP}$ in $J_0(N)/(I_f \cap I_g)J_0(N)$. Note that there is some similarity between these hypotheses and the hypothesis in Proposition 1.3 that q does not divide $|(J_0(N)/F^\vee)(K)_{\text{tor}}|$. In any case, our discussion just above emphasizes the need to study the torsion in $J_0(N)$ and its quotients over number fields other than $\mathbf{Q}$.

The first step in the proof of Theorem 1.4 is the following proposition (also proved in Section 2), which may be of independent interest:

Proposition 1.6. *Let f be an eigenform of weight 2 on $\Gamma_0(N)$ of analytic rank one (i.e., whose L -function vanishes to order one at $s = 1$). Let J' be a quotient of $J = J_0(N)$ through which the map $J \rightarrow A_f$ factors. Let π' denote the map $J' \rightarrow A_f$ and π'' the map $J \rightarrow J'$ in this factorization. Let F' denote the kernel of π' . Thus we have the following diagram:*

$$\begin{array}{ccccccc} & & J & & & & \\ & & \downarrow \pi'' & \searrow \pi & & & \\ 0 & \longrightarrow & F' & \longrightarrow & J' & \xrightarrow{\pi'} & A_f \longrightarrow 0 \end{array}$$

Then

$$|A_f(K)/\pi(\mathbf{TP})| = \left| \frac{J'(K)}{F'(K) + \pi''(\mathbf{TP})} \right| \cdot |\ker(H^1(K, F') \rightarrow H^1(K, J'))|. \quad (2)$$

2 Proofs

In this section, we prove Proposition 1.6 and Theorem 1.4. We start by proving Proposition 1.6.

Proof of Proposition 1.6. Consider the exact sequence $0 \rightarrow F' \rightarrow J' \rightarrow A_f \rightarrow 0$. Part of the associated long exact sequence of Galois cohomology is

$$0 \rightarrow F'(K) \rightarrow J'(K) \xrightarrow{\pi'} A_f(K) \xrightarrow{\delta} H^1(K, F') \rightarrow H^1(K, J') \rightarrow \cdots, \quad (3)$$

where δ denotes the boundary map. Note that in this proof, the letters π' and π'' denote π' and π'' restricted to the K -valued points in their domain.

Since $\pi''(\mathbf{TP}) \subseteq J'(K)$, by the exactness of (3) we see that $\delta(\pi'(\pi''(\mathbf{TP}))) = 0$. Using the exactness of (3) again, we see that δ thus induces a surjection

$$\phi : A_f(K)/\pi'(\pi''(\mathbf{TP})) \rightarrow \ker(H^1(K, F') \rightarrow H^1(K, J')).$$

Since $\pi'(J'(K)) \subseteq \ker(\delta)$, we see that π' induces a natural map $\psi : J'(K) \rightarrow \ker(\phi)$.

Claim: ψ is surjective and its kernel is $F'(K) + \pi''(\mathbf{TP})$.

Proof. Let $x \in J'(K)$. Then

$$\begin{aligned} x \in \ker(\psi) &\iff \pi'(x) = 0 \in \ker(\phi) \hookrightarrow A_f(K)/\pi'(\pi''(\mathbf{TP})) \\ &\iff \pi'(x) \in \pi'(\pi''(\mathbf{TP})) \\ &\iff \exists t \in \mathbf{T} : x - \pi''(tP) \in \ker(\pi') = F'(K) \\ &\iff x \in F'(K) + \pi''(\mathbf{TP}). \end{aligned}$$

Thus $\ker(\psi) = F'(K) + \pi''(\mathbf{TP})$. To prove surjectivity of ψ , note that given an element of $\ker(\phi)$, we can write the element as $y + \pi'(\pi''(\mathbf{TP}))$ for some $y \in A_f(K)$ such that $\delta(y) = 0$. Then by the exactness of (3), $y \in \text{Im}(\pi')$, hence $y + \pi'(\pi''(\mathbf{TP})) \in \text{Im}(\psi)$. $\square$

By the discussion above, we get an exact sequence:

$$0 \rightarrow \frac{J'(K)}{F'(K) + \pi''(\mathbf{TP})} \xrightarrow{\psi'} \frac{A_f(K)}{\pi'(\pi''(\mathbf{TP}))} \xrightarrow{\phi} \ker(H^1(K, F') \rightarrow H^1(K, J')) \rightarrow 0, \quad (4)$$

where ψ' is the natural map induced by ψ . Now

$$|A_f(K)/\pi'(\pi''(\mathbf{TP}))| = |A_f(K)/\pi(\mathbf{TP})|,$$

and the latter group is finite in our situation. Hence all groups in (4) are finite, and Proposition 1.6 now follows from the exactness of (4). $\square$

In the rest of this section, we prove Theorem 1.4. We work in slightly more generality in the beginning and assume that f and g are any newforms (whose Fourier coefficients need not be integers). Thus the associated newform quotients A_f and A_g need not be elliptic curves, but we will still denote them by E and F (respectively) for simplicity of notation.

Recall that $I_g = \text{Ann}_{\mathbf{T}} g$, and let $J' = J/(I_f \cap I_g)J$. The quotient map $\pi : J \rightarrow J/I_f J$ factors through $J' = J/(I_f \cap I_g)J$. By proposition (1.6), it suffices to show that r divides the right side of (2) with our choice of J' , which is what we will do.

Recall that π' denotes the projection $J' \rightarrow A_f = E$, $F' = \ker \pi'$, and π'' denotes the projection $J \rightarrow J'$. Let B denote the kernel of $\pi : J \rightarrow E$, which is the abelian subvariety $I_f J$ of J . We have the following diagram, in which the two sequences of four arrows are exact (one horizontal and one upwards diagonal):

$$\begin{array}{ccccccc}
 & & F^\vee & & E^\vee & & 0 \\
 & & \downarrow \hookrightarrow & & \downarrow \sim & & \nearrow \\
 0 & \longrightarrow & B & \longrightarrow & J & \xrightarrow{\pi} & E \longrightarrow 0 \\
 & & \downarrow & & \downarrow \pi'' & \nearrow \pi' & \\
 & & F' & & J' & & F \\
 & \nearrow & & & \searrow & & \\
 0 & & & & & &
 \end{array}$$

Now F' is connected, since it is a quotient of B (as a simple diagram chase above shows) and B is connected. Thus, by looking at dimensions, one sees that F' is the image of $F^\vee$ under π'' . Since the composite $F^\vee \hookrightarrow J \rightarrow J' \rightarrow F$ is an isogeny, the quotient map $J' \rightarrow F$ induces an isogeny $\pi''(F^\vee) \sim F$, and hence an isogeny $F' \sim F$. Thus F' and F have the same rank (over $\mathbf{Q}$ or over K). Let E' denote $\pi''(E^\vee)$. Since π induces an isogeny from $E^\vee$ to E , we see that π' also induces an isogeny from E' to E . Thus E' and E have the same rank (over $\mathbf{Q}$ or over K).

Now we impose the assumption that f and g have integer Fourier coefficients, so that E and F are elliptic curves. Recall that r is an integer such that (a) f and g are congruent modulo r and for every prime p that divides r , f and g are not congruent modulo a power of p greater than $p^{\text{ord}_p r}$, (b) all maximal ideals $\mathfrak{m}$ of $\mathbf{T}$ whose residue characteristic divides r and that are in the support of $J[I_f]$ or of $J[I_g]$ satisfy multiplicity one, and (c) r is coprime to the order of the torsion subgroup of the projection of $\mathbf{TP}$ in $J/(I_f \cap I_g)J$. Our goal is to show that r divides $|E(K)/\mathbf{Z}\pi(P)|$.

From Lemma 1.2 and on applying π'' , we see that $E'[r] = F'[r]$ and both are direct summands of $E' \cap F'$ as $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ -modules. In particular, the natural maps $H^1(K, E'[r]) \rightarrow H^1(K, E' \cap F')$ and $H^1(K, F'[r]) \rightarrow H^1(K, E' \cap F')$ are injections. Recall that E has analytic rank one and F has Mordell-Weil rank more than one. Then the abelian group $F(K)$ has

rank more than one, and as remarked just before Conjecture 1, the abelian group $E(K)$ has rank one. Also, note that the newform g has analytic rank greater than one, since otherwise the Mordell-Weil rank of F would be at most one. With an eye towards potential generalizations, we remark that after this paragraph, we will not explicitly use the fact that E and F have dimension one (i.e., are elliptic curves). Thus if the conclusions of this paragraph are satisfied, then the rest of the argument would go through even if E and F have dimension greater than one.

Consider the following commutative diagram, where the top and bottom rows are the Kummer exact sequences of E' and F' respectively, and the other maps are the obvious natural maps:

$$\begin{array}{ccccccc}
0 & \longrightarrow & E'(K)/rE'(K) & \longrightarrow & H^1(K, E'[r]) & \longrightarrow & H^1(K, E')[r] \longrightarrow 0 \\
& & & & \parallel & & \searrow \\
& & & & & & H^1(K, E') \\
& & & & \searrow & \nearrow & \\
& & & & & H^1(K, E' \cap F') & \\
& & & & \nearrow & \searrow & \\
& & & & & & H^1(K, F') \\
& & & & \parallel & & \nearrow \\
0 & \longrightarrow & F'(K)/rF'(K) & \xrightarrow{\delta'} & H^1(K, F'[r]) & \longrightarrow & H^1(K, F')[r] \longrightarrow 0
\end{array}$$

Let Q be a generator for the free part of $E'(K)$ (which is isomorphic to the free part of $E(K)$). Then from the top exact sequence in the diagram above, we see that Q gives rise to a non-trivial element σ in $H^1(K, E'[r])$.

Let r' be the smallest positive integer such that $r'\sigma \in \delta'(F'(K)/rF'(K))$ (where δ' is the boundary map in the Kummer exact sequence associated to F' , as indicated in the diagram above). Thus r' divides r (since $r\sigma = 0 \in \delta'(F'(K)/rF'(K))$). Then, by the top and bottom exact sequences in the diagram above, $r'\sigma$ maps to the trivial element in both $H^1(K, E')[r]$ and $H^1(K, F')[r]$, and hence in $H^1(K, E')$ and $H^1(K, F')$. The image of $r'\sigma$ in $H^1(K, E' \cap F')$ is then a non-trivial element of order r/r' that dies in $H^1(K, F')$ and in $H^1(K, E')$. Thus we see that r/r' divides the order of $\ker(H^1(K, E' \cap F') \rightarrow H^1(K, E' \oplus F'))$.

Lemma 2.1. *We have*

$$\frac{J'(K)}{F'(K) + E'(K)} \cong \ker (H^1(K, E' \cap F') \rightarrow H^1(K, E' \oplus F')).$$

Proof. Following a similar situation in [CM00], consider the short exact sequence

$$0 \rightarrow E' \cap F' \rightarrow E' \oplus F' \rightarrow J' \rightarrow 0, \quad (5)$$

where the map $E' \cap F' \rightarrow E' \oplus F'$ is the anti-diagonal embedding $x \mapsto (-x, x)$ and the map $E' \oplus F' \rightarrow J'$ is given by $(x, y) \mapsto x + y$. Part of the associated long exact sequence is

$$\cdots \rightarrow E'(K) \oplus F'(K) \rightarrow J'(K) \rightarrow H^1(K, E' \cap F') \rightarrow H^1(K, E' \oplus F') \rightarrow \cdots,$$

from which we get the lemma. $\square$

By the lemma and the discussion preceding it, we see that r/r' divides $|\frac{J'(K)}{F'(K) + E'(K)}|$. If h is an eigenform of weight 2 on $\Gamma_0(N)$, then $\mathbf{TP} \cap A_h^\vee(K)$ is infinite if and only if h has analytic rank one (this follows by [GZ86, Thm 6.3] if h has analytic rank bigger than one, and the fact that $A_h^\vee(K)$ is finite if h has analytic rank one, by [KL89]). Thus, considering that J' is isogenous to $E' \oplus F'$ and g has analytic rank greater than one, we see that the free parts of $E'(K)$ and of $\pi''(\mathbf{TP})$ agree. Since we are assuming that r is coprime to the order of the torsion part of $\pi''(\mathbf{TP})$ (which is condition (c) on r), considering that r/r' divides $|\frac{J'(K)}{F'(K) + E'(K)}|$, we see that r/r' divides $|\frac{J'(K)}{F'(K) + \pi''(\mathbf{TP})}|$ as well. Thus r/r' divides the first factor on the right side of (2).

If $r' = 1$, then we are done, so let us assume that $r' > 1$. Then $\sigma \notin \delta'(F'(K)/rF'(K))$. So while the image of σ in $H^1(K, E')[r]$ is trivial by the top exact sequence in the diagram above, the image of σ in $H^1(K, F')[r]$ generates a subgroup of order r' , by the lower exact sequence of the diagram above (recall that r' is the *smallest* positive integer such that $r'\sigma \in \delta'(F'(K)/rF'(K))$). Thus, from the same diagram, we see that σ , viewed as an element of $H^1(K, E' \cap F')$, maps to the trivial element of $H^1(K, E')$ but a nontrivial element σ' of $H^1(K, F')$ of order r' . Following a similar situation in [Maz99], considering the exactness of

$$H^1(K, E' \cap F') \rightarrow H^1(K, E') \oplus H^1(K, F') \rightarrow H^1(K, J'),$$

which is part of the long exact sequence associated to (5), we see that the element $(0, \sigma')$ in the middle group dies in the rightmost group, since it

arises from the element σ in the leftmost group. Thus $\sigma' \in H^1(K, F')$ dies in $H^1(K, J')$, and thus is a nontrivial element of order r' of $\ker(H^1(K, F') \rightarrow H^1(K, J'))$. Hence r' divides the second factor on the right side of (2).

Thus r/r' divides the first factor on the right side of (2), and r' divides the second factor on the right side of (2), and so r divides the right side of (2), as was to be shown.

References

- [Aga07] A. Agashe, *A visible factor of the special L-value*, submitted (2007), available at <http://www.math.fsu.edu/~agashe/math.html>.
- [ARS07] A. Agashe, K. Ribet, and W. A. Stein, *The modular degree, congruence primes, and multiplicity one*, preprint (2007), available at <http://www.math.fsu.edu/~agashe/moddeg3.html>.
- [AS02a] A. Agashe and W. A. Stein, Appendix to Joan-C. Lario and René Schoof, *Some computations with Hecke rings and deformation rings*, Experiment. Math. **11** (2002), no. 2, 303–311.
- [AS02b] Amod Agashe and William Stein, *Visibility of Shafarevich-Tate groups of abelian varieties*, J. Number Theory **97** (2002), no. 1, 171–185.
- [AS05] ———, *Visible evidence for the Birch and Swinnerton-Dyer conjecture for modular abelian varieties of analytic rank zero*, Math. Comp. **74** (2005), no. 249, 455–484 (electronic), With an appendix by J. Cremona and B. Mazur.
- [AU96] Ahmed Abbes and Emmanuel Ullmo, *À propos de la conjecture de Manin pour les courbes elliptiques modulaires*, Compositio Math. **103** (1996), no. 3, 269–286. MR 97f:11038
- [CM00] J.E. Cremona and B. Mazur, *Visualizing elements in the Shafarevich-Tate group*, Experiment. Math. **9** (2000), no. 1, 13–28. MR 1 758 797
- [Cre] J.E. Cremona, *Elliptic curve data*, <http://www.warwick.ac.uk/~masgaj/ftp/data/index.html>.

- [DSW03] Neil Dummigan, William Stein, and Mark Watkins, *Constructing elements in Shafarevich-Tate groups of modular motives*, Number theory and algebraic geometry, London Math. Soc. Lecture Note Ser., vol. 303, Cambridge Univ. Press, Cambridge, 2003, pp. 91–118. MR MR2053457
- [Eme03] Matthew Emerton, *Optimal quotients of modular Jacobians*, Math. Ann. **327** (2003), no. 3, 429–458.
- [Gro91] B.H. Gross, *Kolyvagin’s work on modular elliptic curves, L-functions and arithmetic* (Durham, 1989), Cambridge Univ. Press, Cambridge, 1991, pp. 235–256.
- [GZ86] B. Gross and D. Zagier, *Heegner points and derivatives of L-series*, Invent. Math. **84** (1986), no. 2, 225–320. MR 87j:11057
- [KL89] V.A. Kolyvagin and D.Y. Logachev, *Finiteness of the Shafarevich-Tate group and the group of rational points for some modular abelian varieties*, Algebra i Analiz **1** (1989), no. 5, 171–196.
- [Kol90] V. A. Kolyvagin, *Euler systems*, The Grothendieck Festschrift, Vol. II, Progr. Math., vol. 87, Birkhäuser Boston, Boston, MA, 1990, pp. 435–483.
- [Maz77] B. Mazur, *Modular curves and the Eisenstein ideal*, Inst. Hautes Études Sci. Publ. Math. (1977), no. 47, 33–186 (1978).
- [Maz78] ———, *Rational isogenies of prime degree (with an appendix by D. Goldfeld)*, Invent. Math. **44** (1978), no. 2, 129–162.
- [Maz99] ———, *Visualizing elements of order three in the Shafarevich-Tate group*, Asian J. Math. **3** (1999), no. 1, 221–232. MR 2000g:11048
- [Rib90] K.A. Ribet, *On modular representations of $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ arising from modular forms*, Invent. Math. **100** (1990), no. 2, 431–476.
- [Wal85] J.-L. Waldspurger, *Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie*, Compositio Math. **54** (1985), no. 2, 173–242. MR 87g:11061b
- [Wil95] A.J. Wiles, *Modular elliptic curves and Fermat’s last theorem*, Ann. of Math. (2) **141** (1995), no. 3, 443–551.